

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Di- μ -iodido-bis{[(4-fluorobenzoylmethyl-ene)triphenyl- λ^5 -phosphorane]iodido-mercury(II)}

Mehmet Akkurt,^{a*} Kazem Karami,^b Şerife Pınar Yalçın^a and Orhan Büyükgüngör^c

^aDepartment of Physics, Faculty of Arts and Sciences, Erciyes University, 38039 Kayseri, Turkey, ^bDepartment of Chemistry, Isfahan University of Technology, Isfahan, 84156/83111, Iran, and ^cDepartment of Physics, Faculty of Arts and Sciences, Ondokuz Mayıs University, 55139 Samsun, Turkey
Correspondence e-mail: akkurt@erciyes.edu.tr


Received 21 March 2008; accepted 31 March 2008

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.012$ Å; R factor = 0.045; wR factor = 0.090; data-to-parameter ratio = 19.2.

In the title complex, $[\text{Hg}_2\text{I}_4(\text{C}_{26}\text{H}_{20}\text{FOP})_2]$, the Hg^{II} centre is four-coordinate with one short $\text{Hg}-\text{I}$ bond [2.6895 (7) Å], one $\text{Hg}-\text{C}$ bond and two asymmetric bridging $\text{Hg}-\text{I}$ bonds with distances of 2.7780 (8) and 3.2599 (8) Å. The title molecule has a crystallographic inversion centre at the centroid of the four-membered ring formed by the two Hg atoms and two I atoms. The crystal packing is stabilized by $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds.

Related literature

For related literature, see: Baenziger *et al.* (1978); Belluco *et al.* (1996); Bent (1961); Holy *et al.* (1976); Kalyanasundari *et al.* (1995, 1999); Karami (2007); Laavanya *et al.* (2001); Uson *et al.* (1985).



Experimental

Crystal data

$[\text{Hg}_2\text{I}_4(\text{C}_{26}\text{H}_{20}\text{FOP})_2]$

$M_r = 1705.56$

Triclinic, $P\bar{1}$

$a = 10.0346$ (16) Å

$b = 11.8594$ (19) Å

$c = 13.235$ (2) Å

$\alpha = 92.513$ (13)°

$\beta = 111.293$ (12)°

$\gamma = 113.117$ (12)°

$V = 1317.4$ (4) Å³

$Z = 1$

Mo $K\alpha$ radiation

$\mu = 8.27$ mm⁻¹

$T = 293$ (2) K

$0.26 \times 0.17 \times 0.08$ mm

Data collection

Stoe IPDSII diffractometer

Absorption correction: integration

(*X-RED32*; Stoe & Cie, 2002)

$T_{\text{min}} = 0.222$, $T_{\text{max}} = 0.558$

16225 measured reflections

5553 independent reflections

4486 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.156$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.045$

$wR(F^2) = 0.089$

$S = 1.05$

5553 reflections

289 parameters

H-atom parameters constrained

$\Delta\rho_{\text{max}} = 1.00$ e Å⁻³

$\Delta\rho_{\text{min}} = -0.67$ e Å⁻³

Table 1

Selected geometric parameters (Å, °).

Hg1—I1	2.7780 (8)	Hg1—C19	2.281 (5)
Hg1—I2	2.6895 (7)	Hg1—I1 ⁱ	3.2599 (8)
I1—Hg1—I2	111.82 (2)	I1 ⁱ —Hg1—I2	97.77 (2)
I1—Hg1—C19	116.49 (16)	I1 ⁱ —Hg1—C19	96.90 (16)
I1—Hg1—I1 ⁱ	94.17 (2)	Hg1—I1—Hg1 ⁱ	85.84 (2)
I2—Hg1—C19	127.98 (15)		

Symmetry code: (i) $-x + 1, -y + 1, -z + 1$.

Table 2

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C4—H4 \cdots O1 ⁱⁱ	0.93	2.59	3.271 (11)	131
C12—H12 \cdots O1	0.93	2.32	3.124 (8)	144
C22—H22 \cdots O1	0.93	2.44	2.749 (10)	100

Symmetry code: (ii) $x + 1, y, z$.

Data collection: *X-Area* (Stoe & Cie, 2002); cell refinement: *X-Area*; data reduction: *X-RED32* (Stoe & Cie, 2002); program(s) used to solve structure: *SIR97* (Altomare *et al.*, 1999); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

The authors acknowledge the Faculty of Arts and Sciences, Ondokuz Mayıs University, Turkey, for the use of the Stoe IPDSII diffractometer (purchased under grant F. 279 of the University Research Fund).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT2690).

References

- Altomare, A., Burla, M. C., Camalli, M., Cascarano, G. L., Giacovazzo, C., Guagliardi, A., Moliterni, A. G. G., Polidori, G. & Spagna, R. (1999). *J. Appl. Cryst.* **32**, 115–119.
- Baenziger, N. C., Flynn, R. M., Swenson, D. C. & Holy, N. L. (1978). *Acta Cryst.* **B34**, 2300–2301.
- Belluco, U., Michelin, R. A., Bertini, R., Facchin, G., Pace, G., Zanutto, L., Mozzon, M., Furlan, M. & Zangrando, E. (1996). *Inorg. Chim. Acta*, **252**, 355–366.
- Bent, H. A. (1961). *Chem. Rev.* **61**, 275–311.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Holy, N. L., Baenziger, N. C., Flynn, R. M. & Swenson, D. C. (1976). *J. Am. Chem. Soc.* **98**, 7823–7824.

- Kalyanasundari, M., Panchanatheswaran, K., Parthasarathi, V. & Robinson, W. T. (1999). *Bull. Chem. Soc. Jpn*, **72**, 33–36.
- Kalyanasundari, M., Panchanatheswaran, K., Robinson, W. T. & Wen, H. (1995). *J. Organomet. Chem.* **491**, 103–109.
- Karami, K. (2007). *J. Chem. Res. (S)*, pp. 725–727.
- Laavanya, P., Venkatasubramanian, U., Panchanatheswaran, K. & Bauer, J. A. K. (2001). *Chem. Commun.* pp. 1660–1661.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Stoe & Cie (2002). *X-AREA* and *X-RED32*. Stoe & Cie, Darmstadt, Germany.
- Uson, R., Fornies, J., Navarro, R., Espinet, P. & Mendivil, C. (1985). *J. Organomet. Chem.* **290**, 125–131.

supplementary materials

Acta Cryst. (2008). E64, m612-m613 [doi:10.1107/S1600536808008611]

Di- μ -iodido-bis{[(4-fluorobenzoylmethylene)triphenyl- λ^5 -phosphorane]iodidomercury(II)}

M. Akkurt, K. Karami, S.P. Yalçın and O. Büyükgüngör

Comment

The dimeric structure adopted by complexes is in contrast to the O-coordinated trinuclear mercury (II) complex of the phosphorus ylide $\text{Ph}_3\text{PCHCOPh}$ (Kalyanasundari *et al.*, 1999), but is similar to the structure of *trans*-di-iododiodobis (triphenyl phosphoniumcyclopentadienylide) dimercury(II) reported by Baenziger *et al.* (Baenziger *et al.*, 1978) and the C-coordinated dinuclear mercury(II) halide complexes of $\text{Ph}_3\text{CHCOPh}(\text{BPPY})$ (Kalyanasundari *et al.*, 1995). The C-coordination of FBPPY is in stark contrast to the O-coordination of the phosphorus ylide, $\text{Ph}_3\text{PC}(\text{COMe})(\text{COPh})$ (ABPPY), to a Hg^{II} centre (Laavanya *et al.*, 2001). The difference in coordination mode between ABPPY and FBPPY to Hg^{II} can be rationalized in terms of the electronic properties and steric requirements of the ylides. The nucleophilicity of the carbanion in ABPPY is less than for FBPPY; this is due to the additional delocalization of the ylide electron density in ABPPY which is facilitated by the second carbonyl group. This will reduce the ability of ABPPY to bind *via* the ylidic carbon. Belluco *et al.* have studied steric influences on the coordination modes of ylide molecules to Pt(II) systems (Belluco *et al.*, 1996). These authors concluded that the preferred coordination mode is *via* the ylidic carbon, but that steric hindrance around the metal centre or the ylidic carbon will necessitate O-coordination. Indeed, this trend is reflected here, both BPPY and FBPPY are slightly less sterically demanding than ABPPY, and both are C-coordinated to Hg^{II} .

The title molecule has a crystallographic inversion symmetry in the mid-point of the four-membered ring formed by the two Hg atoms and two I atoms (Fig.1). The crystal structure of the title complex reveals that the Hg^{II} centre forms four close contacts with sp^3 hybridization and has a 4-coordinate environment with one short Hg—I bond 2.6895 (7), one Hg—C bond and two asymmetric bridging Hg—I bonds at distances of 2.7780 (8) and 3.2599 (8) Å in complex $[\{\text{HgI}_2(\text{FBPPY})\}_2]$. The significant shortening of the Hg—C bond length, 2.281 (5) Å compared to analogous distances in $[(\text{C}_6\text{H}_5)_3\text{PCHCOC}_6\text{H}_5\text{HgI}_2]_2$ (Kalyanasundari *et al.*, 1995) and in $[(\text{C}_5\text{H}_4\text{P}(\text{C}_6\text{H}_5)_3\text{HgI}_2)_2]$ (Holy *et al.*, 1976) [2.312 (13) and 2.292 (8) Å, respectively] must be attributed to the use of mercury orbitals with high s character for bonding to the ylidic carbon. The use of non-equivalent hybrid orbitals with high s character to bond to low electronegative atoms was proposed by Bent in the concept of isovalent hybridization to account for the variation in bond lengths and bond angles around a central atom (Bent, 1961). The terminal Hg—I bond length, 2.7780 (8) Å is comparable to 2.615 Å observed in the case of $\text{Hg}_2\text{I}_4(\text{ABPPY})_2$, which has a tetrahedral coordination environment around mercury with a bridging structure (Laavanya *et al.*, 2001). The two bridged Hg—I bonds fall within the range 2.778 - 3.25994 Å reported for other structures (Laavanya *et al.*, 2001) containing chloro bridged mercury. The angles around mercury vary from 94.17 (2) to 111.82 (2) for I—Hg—I, a very distorted tetrahedral environment. This distortion must be due to the higher s character of the sp^3 hybrid mercury orbitals involved in the above bonds and the formation of a strong Iodo bridge between the Hg atoms which requires the internal I—Hg—I angle to be considerably smaller. The stabilized resonance structure for the title ylide is destroyed by the complexes formation. On the other hand, the bond length of P(1)—C(19) in the similar ylide is 1.706 Å (Uson *et al.*, 1985) which shows that the above bond is considerably elongated to 1.787 (6) Å in complex $[\{\text{HgI}_2(\text{FBPPY})\}_2]$. The adaptation

supplementary materials

of dimeric structure in Hg^{II} ylide complex may be explained by both the preference of Hg^{II} to four coordination and the stability of the 18 electron configuration around Hg^{II} .

Experimental

To a chloroform solution (15 ml) of triphenylphosphine (1 mmol) was added 2-bromo-4-fluoroacetophenone (1 mmol) and the resulting mixture was stirred for 12 h. The solution was filtered off, and the precipitate washed with diethyl-ether and air-dried. Further treatment with aqueous NaOH solution (0.5M) led to elimination of HBr, giving the free ligand precursors FBPPY. To a solution of FBPPY (0.100 g, 0.25 mmol) in acetone (5 ml) was added mercury (II) iodide (0.114 g, 0.25 mmol). The mixture was stirred for 12 h. On concentration by removing the solvent by vacuum, a pale yellow precipitate was obtained. The products were washed with benzene and dried *in vacuo*. Yield: 81%, *M.p.* 214 °C. Analysis calculated for $\text{C}_{52}\text{H}_{40}\text{F}_2\text{Hg}_2\text{I}_4\text{O}_2\text{P}_2 \cdot \text{C}$ C 36.6, H 2.4%. Found: C 36.45, H 2.3%, ^1H NMR: 4.62(d, 1H, CH, $2J_{\text{PH}} = 5.5$ Hz), 7.1–8 (m, 19H, Ph) p.p.m. and ^{31}P NMR: 20.34 p.p.m. (Karami, 2007).

Refinement

H atoms were placed in calculated positions and refined using a riding model with $\text{C—H} = 0.93$ Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. An ORTEP-3 view of the title compound with the atom numbering scheme. Displacement ellipsoids for non-H atoms are drawn at the 30% probability level.




Fig. 2. The packing and intermolecular hydrogen bonding interactions of the title compound.

Di- μ -iodido- bis{[(4-fluorobenzoylmethylene)triphenyl- λ^5 -phosphorane]iodidomercury(II)}

Crystal data

$[\text{Hg}_2\text{I}_4(\text{C}_{26}\text{H}_{20}\text{FOP})_2]$

$M_r = 1705.56$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 10.0346$ (16) Å

$Z = 1$

$F_{000} = 788$

$D_x = 2.150$ Mg m $^{-3}$

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 47681 reflections

$b = 11.8594 (19) \text{ \AA}$
 $c = 13.235 (2) \text{ \AA}$
 $\alpha = 92.513 (13)^\circ$
 $\beta = 111.293 (12)^\circ$
 $\gamma = 113.117 (12)^\circ$
 $V = 1317.4 (4) \text{ \AA}^3$

$\theta = 1.7\text{--}28.3^\circ$
 $\mu = 8.27 \text{ mm}^{-1}$
 $T = 293 (2) \text{ K}$
 Plate, colourless
 $0.26 \times 0.17 \times 0.08 \text{ mm}$

Data collection

Stoe IPDSII diffractometer	5553 independent reflections
Monochromator: plane graphite	4486 reflections with $I > 2\sigma(I)$
Detector resolution: $6.67 \text{ pixels mm}^{-1}$	$R_{\text{int}} = 0.156$
$T = 293(2) \text{ K}$	$\theta_{\text{max}} = 27.0^\circ$
ω scans	$\theta_{\text{min}} = 1.7^\circ$
Absorption correction: integration (X-RED32; Stoe & Cie, 2002)	$h = -12 \rightarrow 12$
$T_{\text{min}} = 0.222, T_{\text{max}} = 0.558$	$k = -15 \rightarrow 15$
16225 measured reflections	$l = -16 \rightarrow 16$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.045$	H-atom parameters constrained
$wR(F^2) = 0.089$	$w = 1/[\sigma^2(F_o^2) + (0.0388P)^2 + 0.7953P]$
$S = 1.05$	where $P = (F_o^2 + 2F_c^2)/3$
5553 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
289 parameters	$\Delta\rho_{\text{max}} = 1.00 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\text{min}} = -0.67 \text{ e \AA}^{-3}$
	Extinction correction: none

Special details

Geometry. Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement on F^2 for ALL reflections except those flagged by the user for potential systematic errors. Weighted R -factors wR and all goodnesses of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The observed criterion of $F^2 > \sigma(F^2)$ is used only for calculating $-R$ -factor-obs *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

supplementary materials

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Hg1	0.53557 (3)	0.65438 (3)	0.42856 (2)	0.0581 (1)
I1	0.29177 (6)	0.52860 (4)	0.49338 (4)	0.0650 (2)
I2	0.76379 (6)	0.86528 (5)	0.57953 (4)	0.0727 (2)
P1	0.58806 (17)	0.74049 (12)	0.20232 (11)	0.0389 (4)
F1	-0.1102 (8)	0.0461 (5)	0.0631 (6)	0.133 (3)
O1	0.2430 (5)	0.6313 (4)	0.1541 (4)	0.0613 (16)
C1	0.7783 (7)	0.7396 (5)	0.2330 (5)	0.0445 (17)
C2	0.8580 (8)	0.7185 (7)	0.3349 (5)	0.062 (2)
C3	0.9949 (9)	0.7038 (8)	0.3566 (6)	0.069 (3)
C4	1.0532 (9)	0.7126 (8)	0.2760 (7)	0.074 (3)
C5	0.9746 (10)	0.7344 (9)	0.1756 (7)	0.080 (3)
C6	0.8373 (9)	0.7470 (7)	0.1524 (7)	0.065 (3)
C7	0.6204 (7)	0.8894 (5)	0.2717 (5)	0.0442 (17)
C8	0.7676 (8)	0.9929 (6)	0.3076 (6)	0.061 (2)
C9	0.7889 (11)	1.1059 (6)	0.3622 (7)	0.076 (3)
C10	0.6674 (12)	1.1146 (7)	0.3787 (6)	0.078 (3)
C11	0.5234 (11)	1.0153 (7)	0.3426 (7)	0.072 (3)
C12	0.4974 (9)	0.9025 (6)	0.2876 (6)	0.061 (2)
C13	0.4969 (7)	0.7270 (5)	0.0543 (5)	0.0423 (17)
C14	0.5281 (8)	0.8339 (6)	0.0109 (5)	0.0548 (19)
C15	0.4683 (10)	0.8238 (7)	-0.1016 (6)	0.068 (3)
C16	0.3754 (10)	0.7082 (7)	-0.1731 (6)	0.069 (2)
C17	0.3410 (9)	0.6005 (6)	-0.1319 (5)	0.061 (2)
C18	0.4023 (8)	0.6097 (5)	-0.0176 (5)	0.0527 (19)
C19	0.4769 (6)	0.6070 (5)	0.2440 (4)	0.0401 (17)
C20	0.3012 (7)	0.5596 (6)	0.1835 (5)	0.0463 (17)
C21	0.1984 (7)	0.4217 (6)	0.1586 (5)	0.0539 (19)
C22	0.0358 (9)	0.3826 (8)	0.1146 (8)	0.078 (3)
C23	-0.0692 (11)	0.2564 (10)	0.0825 (10)	0.107 (4)
C24	-0.0083 (12)	0.1715 (8)	0.0951 (8)	0.088 (3)
C25	0.1511 (11)	0.2031 (7)	0.1366 (7)	0.081 (3)
C26	0.2542 (8)	0.3310 (6)	0.1691 (6)	0.063 (2)
H2	0.81880	0.71420	0.38910	0.0740*
H3	1.04710	0.68820	0.42450	0.0830*
H4	1.14600	0.70380	0.28970	0.0890*
H5	1.01540	0.74060	0.12210	0.0960*
H6	0.78430	0.76050	0.08360	0.0780*
H8	0.85050	0.98690	0.29550	0.0730*
H9	0.88720	1.17580	0.38750	0.0900*
H10	0.68370	1.19050	0.41570	0.0940*
H11	0.44150	1.02310	0.35490	0.0860*
H12	0.39720	0.83480	0.26100	0.0730*
H14	0.59000	0.91280	0.05830	0.0650*
H15	0.49070	0.89610	-0.13000	0.0820*
H16	0.33570	0.70260	-0.24940	0.0820*

H17	0.27730	0.52210	-0.18020	0.0730*
H18	0.38010	0.53740	0.01070	0.0630*
H19	0.50550	0.53940	0.23070	0.0480*
H22	-0.00370	0.44230	0.10640	0.0940*
H23	-0.17860	0.23040	0.05300	0.1290*
H25	0.18880	0.14240	0.14280	0.0970*
H26	0.36340	0.35610	0.19860	0.0750*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Hg1	0.0633 (2)	0.0652 (2)	0.0484 (1)	0.0285 (1)	0.0256 (1)	0.0120 (1)
I1	0.0709 (3)	0.0725 (3)	0.0830 (3)	0.0449 (2)	0.0483 (3)	0.0334 (2)
I2	0.0717 (3)	0.0778 (3)	0.0580 (3)	0.0308 (2)	0.0200 (2)	-0.0026 (2)
P1	0.0372 (7)	0.0383 (7)	0.0415 (7)	0.0166 (6)	0.0172 (6)	0.0033 (5)
F1	0.110 (4)	0.067 (3)	0.161 (6)	-0.022 (3)	0.062 (4)	-0.009 (3)
O1	0.049 (2)	0.065 (3)	0.074 (3)	0.033 (2)	0.021 (2)	0.015 (2)
C1	0.043 (3)	0.039 (3)	0.045 (3)	0.014 (2)	0.017 (2)	-0.002 (2)
C2	0.050 (4)	0.094 (5)	0.036 (3)	0.036 (3)	0.010 (3)	-0.003 (3)
C3	0.054 (4)	0.094 (5)	0.049 (4)	0.037 (4)	0.009 (3)	-0.005 (3)
C4	0.050 (4)	0.094 (5)	0.079 (5)	0.036 (4)	0.025 (4)	0.003 (4)
C5	0.077 (5)	0.120 (7)	0.084 (5)	0.056 (5)	0.059 (5)	0.041 (5)
C6	0.066 (4)	0.083 (5)	0.070 (4)	0.040 (4)	0.043 (4)	0.030 (3)
C7	0.048 (3)	0.046 (3)	0.044 (3)	0.027 (3)	0.018 (2)	0.008 (2)
C8	0.057 (4)	0.056 (4)	0.055 (4)	0.024 (3)	0.010 (3)	-0.007 (3)
C9	0.078 (5)	0.045 (4)	0.081 (5)	0.019 (3)	0.020 (4)	-0.006 (3)
C10	0.110 (7)	0.061 (4)	0.067 (4)	0.059 (5)	0.019 (4)	-0.001 (3)
C11	0.089 (6)	0.071 (5)	0.074 (5)	0.051 (4)	0.036 (4)	0.007 (4)
C12	0.068 (4)	0.058 (4)	0.070 (4)	0.036 (3)	0.033 (4)	0.011 (3)
C13	0.041 (3)	0.043 (3)	0.046 (3)	0.021 (2)	0.019 (2)	0.005 (2)
C14	0.057 (4)	0.048 (3)	0.049 (3)	0.018 (3)	0.017 (3)	0.007 (2)
C15	0.088 (5)	0.059 (4)	0.059 (4)	0.031 (4)	0.033 (4)	0.022 (3)
C16	0.090 (5)	0.074 (4)	0.042 (3)	0.044 (4)	0.019 (3)	0.009 (3)
C17	0.073 (4)	0.058 (4)	0.045 (3)	0.031 (3)	0.015 (3)	-0.002 (3)
C18	0.060 (4)	0.038 (3)	0.055 (3)	0.020 (3)	0.021 (3)	0.002 (2)
C19	0.040 (3)	0.043 (3)	0.041 (3)	0.019 (2)	0.020 (2)	0.008 (2)
C20	0.041 (3)	0.056 (3)	0.042 (3)	0.022 (3)	0.017 (2)	0.007 (2)
C21	0.045 (3)	0.059 (4)	0.048 (3)	0.011 (3)	0.023 (3)	0.003 (3)
C22	0.048 (4)	0.074 (5)	0.105 (6)	0.017 (4)	0.036 (4)	0.002 (4)
C23	0.052 (5)	0.098 (7)	0.136 (9)	-0.001 (5)	0.043 (5)	-0.018 (6)
C24	0.088 (6)	0.061 (5)	0.082 (5)	-0.008 (5)	0.046 (5)	-0.001 (4)
C25	0.087 (6)	0.052 (4)	0.077 (5)	0.012 (4)	0.025 (4)	0.012 (3)
C26	0.052 (4)	0.054 (4)	0.059 (4)	0.010 (3)	0.014 (3)	0.007 (3)

Geometric parameters (\AA , $^\circ$)

Hg1—I1	2.7780 (8)	C19—C20	1.491 (10)
Hg1—I2	2.6895 (7)	C20—C21	1.493 (9)
Hg1—C19	2.281 (5)	C21—C22	1.382 (13)

supplementary materials

Hg1—I1 ⁱ	3.2599 (8)	C21—C26	1.385 (11)
P1—C1	1.806 (8)	C22—C23	1.383 (15)
P1—C7	1.805 (6)	C23—C24	1.356 (17)
P1—C13	1.805 (6)	C24—C25	1.369 (17)
P1—C19	1.787 (6)	C25—C26	1.394 (11)
F1—C24	1.369 (11)	C2—H2	0.9300
O1—C20	1.212 (9)	C3—H3	0.9300
C1—C2	1.387 (9)	C4—H4	0.9300
C1—C6	1.387 (12)	C5—H5	0.9300
C2—C3	1.381 (13)	C6—H6	0.9300
C3—C4	1.381 (13)	C8—H8	0.9300
C4—C5	1.373 (12)	C9—H9	0.9300
C5—C6	1.369 (15)	C10—H10	0.9300
C7—C8	1.389 (10)	C11—H11	0.9300
C7—C12	1.389 (13)	C12—H12	0.9300
C8—C9	1.394 (11)	C14—H14	0.9300
C9—C10	1.355 (17)	C15—H15	0.9300
C10—C11	1.348 (14)	C16—H16	0.9300
C11—C12	1.375 (11)	C17—H17	0.9300
C13—C14	1.385 (9)	C18—H18	0.9300
C13—C18	1.390 (8)	C19—H19	0.9800
C14—C15	1.368 (9)	C22—H22	0.9300
C15—C16	1.373 (11)	C23—H23	0.9300
C16—C17	1.378 (10)	C25—H25	0.9300
C17—C18	1.391 (9)	C26—H26	0.9300
Hg1…C2	3.694 (9)	C15…H23 ^x	3.0300
Hg1…C12	3.624 (7)	C19…H2	2.9400
Hg1…C26	4.216 (7)	C19…H18	2.8500
Hg1…H2	2.8900	C19…H26	2.6800
Hg1…H12	3.4500	C20…H12	3.0300
Hg1…H26	3.9000	C20…H18	2.7100
I2…C25 ⁱ	3.739 (8)	C23…H5 ^{ix}	2.9500
I2…C7	3.863 (6)	C24…H5 ^{ix}	3.0900
I1…H2 ⁱ	3.3300	C26…H19	2.5600
I1…H10 ⁱⁱ	3.3800	H2…Hg1	2.8900
I2…H2	3.3600	H2…I2	3.3600
I2…H8 ⁱⁱⁱ	3.2500	H2…C19	2.9400
I2…H11 ⁱⁱ	3.1800	H2…I1 ⁱ	3.3300
F1…C14 ^{iv}	3.292 (11)	H3…C9 ⁱⁱⁱ	3.0800
F1…H14 ^{iv}	2.7700	H3…C10 ⁱⁱⁱ	3.0400
O1…C4 ^v	3.271 (11)	H3…H10 ⁱⁱⁱ	2.5200
O1…C12	3.124 (8)	H4…O1 ^{vi}	2.5900
O1…C13	3.135 (9)	H4…H12 ^{vi}	2.5700
O1…C18	3.270 (9)	H5…C23 ^{ix}	2.9500
O1…H4 ^v	2.5900	H5…C24 ^{ix}	3.0900
O1…H12	2.3200	H6…C13	2.6300

O1...H22	2.4400	H6...C14	2.8900
C2...Hg1	3.694 (9)	H8...C1	2.7400
C4...O1 ^{vi}	3.271 (11)	H8...I2 ⁱⁱⁱ	3.2500
C6...C14	3.558 (12)	H10...I1 ⁱⁱ	3.3800
C7...I2	3.863 (6)	H10...H3 ⁱⁱⁱ	2.5200
C10...C16 ^{vii}	3.517 (11)	H10...H16 ^{vii}	2.5700
C12...O1	3.124 (8)	H11...I2 ⁱⁱ	3.1800
C12...Hg1	3.624 (7)	H12...Hg1	3.4500
C13...O1	3.135 (9)	H12...O1	2.3200
C14...C6	3.558 (12)	H12...C20	3.0300
C14...F1 ^{viii}	3.292 (11)	H12...H4 ^v	2.5700
C16...C10 ^{vii}	3.517 (11)	H14...F1 ^{viii}	2.7700
C18...C20	3.177 (10)	H14...C7	2.7700
C18...O1	3.270 (9)	H14...C8	3.0200
C20...C18	3.177 (10)	H15...C10 ^{vii}	3.0600
C25...I2 ⁱ	3.739 (8)	H15...C11 ^{vii}	3.0200
C26...Hg1	4.216 (7)	H16...C10 ^{vii}	2.8200
C1...H17 ^{ix}	2.9300	H16...H10 ^{vii}	2.5700
C1...H8	2.7400	H17...C1 ^{ix}	2.9300
C2...H19	3.0400	H17...C2 ^{ix}	2.9300
C2...H17 ^{ix}	2.9300	H17...C3 ^{ix}	3.0300
C3...H17 ^{ix}	3.0300	H17...C6 ^{ix}	3.0300
C6...H17 ^{ix}	3.0300	H18...C19	2.8500
C7...H14	2.7700	H18...C20	2.7100
C8...H14	3.0200	H19...C2	3.0400
C9...H3 ⁱⁱⁱ	3.0800	H19...C26	2.5600
C10...H15 ^{vii}	3.0600	H19...H26	2.0000
C10...H16 ^{vii}	2.8200	H22...O1	2.4400
C10...H3 ⁱⁱⁱ	3.0400	H23...C14 ^x	3.0500
C11...H15 ^{vii}	3.0200	H23...C15 ^x	3.0300
C13...H6	2.6300	H26...Hg1	3.9000
C14...H23 ^x	3.0500	H26...C19	2.6800
C14...H6	2.8900	H26...H19	2.0000
I1—Hg1—I2	111.82 (2)	C22—C23—C24	118.2 (11)
I1—Hg1—C19	116.49 (16)	F1—C24—C23	119.1 (11)
I1—Hg1—I1 ⁱ	94.17 (2)	F1—C24—C25	116.9 (9)
I2—Hg1—C19	127.98 (15)	C23—C24—C25	124.0 (9)
I1 ⁱ —Hg1—I2	97.77 (2)	C24—C25—C26	116.6 (9)
I1 ⁱ —Hg1—C19	96.90 (16)	C21—C26—C25	121.8 (8)
Hg1—I1—Hg1 ⁱ	85.84 (2)	C1—C2—H2	120.00
C1—P1—C7	109.1 (3)	C3—C2—H2	120.00
C1—P1—C13	106.6 (3)	C2—C3—H3	121.00
C1—P1—C19	106.1 (3)	C4—C3—H3	120.00

supplementary materials

C7—P1—C13	108.4 (3)	C3—C4—H4	120.00
C7—P1—C19	114.7 (3)	C5—C4—H4	120.00
C13—P1—C19	111.6 (3)	C4—C5—H5	119.00
P1—C1—C2	119.1 (6)	C6—C5—H5	119.00
P1—C1—C6	120.8 (6)	C1—C6—H6	120.00
C2—C1—C6	119.7 (8)	C5—C6—H6	121.00
C1—C2—C3	120.7 (7)	C7—C8—H8	121.00
C2—C3—C4	119.0 (7)	C9—C8—H8	121.00
C3—C4—C5	120.1 (10)	C8—C9—H9	120.00
C4—C5—C6	121.4 (9)	C10—C9—H9	120.00
C1—C6—C5	119.1 (8)	C9—C10—H10	119.00
P1—C7—C8	120.4 (6)	C11—C10—H10	119.00
P1—C7—C12	120.4 (5)	C10—C11—H11	120.00
C8—C7—C12	119.2 (6)	C12—C11—H11	120.00
C7—C8—C9	118.8 (8)	C7—C12—H12	120.00
C8—C9—C10	120.4 (8)	C11—C12—H12	120.00
C9—C10—C11	121.3 (8)	C13—C14—H14	120.00
C10—C11—C12	120.1 (11)	C15—C14—H14	120.00
C7—C12—C11	120.2 (8)	C14—C15—H15	120.00
P1—C13—C14	120.0 (5)	C16—C15—H15	120.00
P1—C13—C18	120.6 (4)	C15—C16—H16	120.00
C14—C13—C18	119.3 (6)	C17—C16—H16	120.00
C13—C14—C15	120.2 (6)	C16—C17—H17	120.00
C14—C15—C16	120.8 (7)	C18—C17—H17	120.00
C15—C16—C17	120.1 (7)	C13—C18—H18	120.00
C16—C17—C18	119.6 (6)	C17—C18—H18	120.00
C13—C18—C17	120.0 (5)	Hg1—C19—H19	109.00
Hg1—C19—P1	110.7 (3)	P1—C19—H19	109.00
Hg1—C19—C20	106.7 (4)	C20—C19—H19	109.00
P1—C19—C20	113.5 (4)	C21—C22—H22	119.00
O1—C20—C19	120.8 (6)	C23—C22—H22	119.00
O1—C20—C21	120.6 (7)	C22—C23—H23	121.00
C19—C20—C21	118.6 (6)	C24—C23—H23	121.00
C20—C21—C22	117.0 (7)	C24—C25—H25	122.00
C20—C21—C26	124.6 (7)	C26—C25—H25	122.00
C22—C21—C26	118.3 (7)	C21—C26—H26	119.00
C21—C22—C23	121.2 (9)	C25—C26—H26	119.00
I2—Hg1—I1—Hg1 ⁱ	-100.09 (2)	P1—C1—C2—C3	173.0 (6)
C19—Hg1—I1—Hg1 ⁱ	99.83 (17)	C1—C2—C3—C4	1.1 (12)
I1 ⁱ —Hg1—I1—Hg1 ⁱ	0.00 (4)	C2—C3—C4—C5	-0.7 (13)
I1 ⁱ —Hg1 ⁱ —I1—Hg1	0.00 (5)	C3—C4—C5—C6	-0.5 (14)
I2 ⁱ —Hg1 ⁱ —I1—Hg1	-112.71 (2)	C4—C5—C6—C1	1.1 (13)
C19 ⁱ —Hg1 ⁱ —I1—Hg1	117.34 (16)	C8—C7—C12—C11	-2.8 (10)
I2—Hg1—C19—C20	-131.8 (3)	P1—C7—C8—C9	-179.1 (6)
I1 ⁱ —Hg1—C19—C20	122.7 (4)	P1—C7—C12—C11	178.6 (6)
I2—Hg1—C19—P1	-7.9 (4)	C12—C7—C8—C9	2.3 (10)
I1 ⁱ —Hg1—C19—P1	-113.4 (3)	C7—C8—C9—C10	-0.7 (11)

I1—Hg1—C19—P1	148.4 (2)	C8—C9—C10—C11	-0.4 (12)
I1—Hg1—C19—C20	24.6 (4)	C9—C10—C11—C12	-0.2 (12)
C19—P1—C1—C2	-45.5 (6)	C10—C11—C12—C7	1.8 (12)
C7—P1—C1—C6	-108.0 (6)	P1—C13—C14—C15	175.4 (7)
C7—P1—C1—C2	78.5 (6)	C18—C13—C14—C15	-1.0 (13)
C13—P1—C1—C2	-164.6 (5)	P1—C13—C18—C17	-175.9 (7)
C1—P1—C7—C12	-162.1 (5)	C14—C13—C18—C17	0.5 (13)
C13—P1—C1—C6	8.8 (6)	C13—C14—C15—C16	0.7 (15)
C19—P1—C1—C6	127.9 (5)	C14—C15—C16—C17	0.1 (16)
C7—P1—C19—C20	84.7 (5)	C15—C16—C17—C18	-0.7 (16)
C13—P1—C19—C20	-39.2 (5)	C16—C17—C18—C13	0.3 (14)
C7—P1—C13—C14	29.7 (8)	Hg1—C19—C20—O1	89.0 (6)
C19—P1—C13—C14	157.0 (6)	Hg1—C19—C20—C21	-91.8 (6)
C1—P1—C13—C18	88.8 (7)	P1—C19—C20—O1	-33.2 (8)
C13—P1—C7—C12	82.2 (6)	P1—C19—C20—C21	146.1 (5)
C19—P1—C7—C12	-43.3 (6)	O1—C20—C21—C22	-8.3 (10)
C1—P1—C19—C20	-154.9 (4)	O1—C20—C21—C26	166.9 (7)
C1—P1—C13—C14	-87.6 (7)	C19—C20—C21—C22	172.4 (7)
C7—P1—C13—C18	-154.0 (7)	C19—C20—C21—C26	-12.4 (10)
C19—P1—C13—C18	-26.7 (8)	C20—C21—C22—C23	176.0 (9)
C1—P1—C7—C8	19.3 (6)	C26—C21—C22—C23	0.5 (13)
C13—P1—C19—Hg1	-159.0 (3)	C20—C21—C26—C25	-175.1 (7)
C13—P1—C7—C8	-96.3 (6)	C22—C21—C26—C25	0.0 (11)
C7—P1—C19—Hg1	-35.2 (4)	C21—C22—C23—C24	-0.2 (16)
C1—P1—C19—Hg1	85.3 (3)	C22—C23—C24—F1	-179.5 (9)
C19—P1—C7—C8	138.1 (5)	C22—C23—C24—C25	-0.8 (17)
C2—C1—C6—C5	-0.6 (11)	F1—C24—C25—C26	-180.0 (8)
P1—C1—C6—C5	-174.0 (6)	C23—C24—C25—C26	1.3 (15)
C6—C1—C2—C3	-0.5 (11)	C24—C25—C26—C21	-0.9 (12)

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $-x+1, -y+2, -z+1$; (iii) $-x+2, -y+2, -z+1$; (iv) $x-1, y-1, z$; (v) $x-1, y, z$; (vi) $x+1, y, z$; (vii) $-x+1, -y+2, -z$; (viii) $x+1, y+1, z$; (ix) $-x+1, -y+1, -z$; (x) $-x, -y+1, -z$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C4—H4 \cdots O1 ^{vi}	0.93	2.59	3.271 (11)	131
C12—H12 \cdots O1	0.93	2.32	3.124 (8)	144
C22—H22 \cdots O1	0.93	2.44	2.749 (10)	100

Symmetry codes: (vi) $x+1, y, z$.

Fig. 1




Fig. 2

